

FUSE SPECTROSCOPY OF THE TWO PROTOTYPE WHITE DWARFS WITH SIGNATURES OF A SUPER-HOT WIND

K. WERNER AND S. DREIZLER

Universität Tübingen, Germany

J.W. KRUK

Johns Hopkins University, U.S.A.

AND

M.L. SITKO

University of Cincinnati, U.S.A.

The O VIII phenomenon describes the occurrence of ultra-high ionization absorption lines of the CNO elements (e.g. O VIII, N VII, C VI, and even Ne X) in the optical spectra hot of DO WDs. This requires temperatures of almost 10^6 K. Since the first discovery of two such objects (Werner et al. 1995) we realized that 50% of all DO WDs are affected (Dreizler et al. 1995), hence it could be that all DO WDs experience this phenomenon at the beginning of the WD cooling sequence. The He II profiles are symmetric, hence line formation occurs in hydrostatic layers. However, we cannot uniquely fit the unusually strong profiles with any model atmosphere, preventing precise parameter determinations. The optical high ionization CNO lines have pronounced blue wings, corresponding to outflow velocities of up to several thousand km/s. We stress that none of the objects was detected in the X-ray region. The inferred low X-ray luminosity confirms that we do not see absorption lines from super-hot photospheres, but that the high temperatures are possibly localized along shock fronts in the wind.

In the UV range (IUE, HST) no high ionization CNO line is detectable, except possibly O VIII 2978Å (Werner et al. 1997, 1999). This negative result is still not understood. On the other hand we see two unidentified broad absorption features at 1420Å and 1500Å. The 1500Å feature may be the blue shifted absorption component from a P Cygni profile of C IV 1550Å. This is corroborated by a very weak and broad emission bump near the rest wavelength. The terminal wind speed is about 10 000 km/s and mass-loss could be driven by radiation pressure. The only photospheric line clearly identified is He II 1640Å. The presence of O V 1371Å is uncertain and the C IV resonance doublet could be of interstellar origin.




Figure 1. Identification of sulfur and phosphorus lines in the FUSE spectra

We performed FUSE observations of the two prototype DOs. HS0713+3958 was recorded on Nov 21, 2000, for 9825 sec and HE0504-2408 on Dec 5, 2001, for 6625 sec. The spectra confirm earlier ORFEUS observations (Werner et al. 1999) showing that the FUV spectral shape is surprisingly flat. Strong reddening by interstellar dust alone is unlikely because no $2200\text{\AA}$ feature is seen ($E(B-V)$ below 0.05). With the superior capabilities of FUSE it was hoped to identify opacity sources responsible for the flat spectral shape, however, we see no hint to any potential absorbers, probably due to insufficient S/N. It is possible, that a dense curtain of weak lines from the iron group elements remains undetected. Note that a similar flat UV shape observed in the sdOB EC11481-2303 results from heavy line blanketing (see Hammer et al. this proceedings). The usual ISM absorption lines can be identified. H_2 is very weak in HE0504-2408 but prominent in HS0713+3958. The only photospheric lines that can be identified in both objects are from S VI and P V (Fig. 1). In addition, HS0713+3958 shows a weak O VI resonance doublet which, however, can also stem from the ISM.

References

- Dreizler S., Heber U., Napiwotzki R., & Hagen H.-J. 1995, A&A 303, L53
Werner K., Dreizler S., Heber U., et al. 1995, A&A 293, L75
Werner K., Dreizler S., Heber U., & Rauch T. 1997, in White Dwarfs, ed. J. Isern, M. Hernanz, E. Garcia-Berro, p. 207
Werner K., Dreizler S., Rauch T., et al. 1999, in White Dwarfs, ed. J.-E. Solheim, E. Meistas, ASP Conf. Series 169, 511